Carbon Nanotubes can orchestrate the immunological response? Perspectives in vaccine application

Elidamar Nunes

Short Communication

Abstract

Although several infections are challenging to treat, advances in nanobiotechnology allowed the utilization of nanotechnology, for example, nanovaccines. Nanotechnology vaccines are powerful in enhancing the immunogenicity over a precise antigen, presenting advantages over other adjuvant approaches, such as expanded stability, prolonged release, decreased immunotoxicity, and immunogenic selectivity.Objective: In this review, we introduce recent advances in carbon nanotubes (CNTs) used in nanovaccines in order to induce either a carrier effect or an immunostimulatory effect. Results: CNTs have the potential to model the immune system, providing, between another properties, their adjuvant effects, besides having an excellent carrier effect. Several studies concerning CNT-based nanotechnology vaccines reveal that the immunogenicity related to the vaccine antigen is increased, due to the immunomodulatory and carrier effects related to CNTs. Also, due to the CNTs properties of being able to carry immunogenic molecules, they can act like non-classical vaccines, which aim to have both a molecule-carrier effect and an immunogenic effect, a quality not found in vaccines with traditional formulations. Adapting and modifying the physicochemical properties regarding CNTs for usage in vaccines may additionally enlarge their efficacy in inducing a T cell-based immune response, fundamental in imitation of the battle against several infections in immunotherapeutic process.

Biography

Elidamar Nunes has expertise on genomic signatures evolution analysis and specific sites and epitopes selection and recognition for development of vaccines efficiency and anti-HIV drugs. Very motivated and interested in various Science projects and research experiments. Currently working on projects that involves the genome viral evolution and nanotechnology to HIV vaccine at University of Sao Paulo and Infectious Diseases Division, Department of Medicine, Federal University of São Paulo in collaboration to Telecommunications and Control Engineering Department, Engineering School, both in Brazil.

Note: - This work is partly presented at joint event on International Conference on Virology and Immunology (July 19-20, 2021 | Paris, France)